

**Instructions:** Show all work. Answers without work required to obtain the solution will not receive full credit. Some questions may contain multiple parts: be sure to answer all of them. Give exact answers unless specifically asked to estimate.


1. Find the volume of the solid under the function  $f(x, y) = 6 - y$  on the region bounded by  $0 \leq x \leq 3, 0 \leq y \leq 4$  using a double integral. (9 points)

$$\int_0^3 \int_0^4 6 - y \, dy \, dx =$$

$$\int_0^3 \left. 6y - \frac{1}{2}y^2 \right|_0^4 dx =$$

$$\int_0^3 24 - 8 \, dx = \int_0^3 16 \, dx =$$

$$16x \Big|_0^3 = \boxed{48}$$




2. The integral  $\int_0^1 \int_{3x}^3 6e^{y^2} dy dx$  can be evaluated only by changing the order of integration. Sketch the region of integration, reverse the order of integration, and evaluate the integral. (8 points)

$$\int_0^3 \int_0^{1/3 y} 6e^{y^2} dx dy =$$

$$\int_0^3 6e^{y^2} \cdot x \Big|_0^{1/3 y} dy =$$

$$\int_0^3 2ye^{y^2} dy = e^{y^2} \Big|_0^3 =$$

$$\boxed{e^9 - 1}$$




3. Evaluate  $\int_0^4 \int_0^{\sqrt{16-x^2}} y dy dx$  by converting to polar coordinates. (8 points)

$$\int_0^{\pi/2} \int_0^4 r \sin \theta r dr d\theta =$$


$$\int_0^{\pi/2} \int_0^4 r^2 \sin \theta dr d\theta = \int_0^{\pi/2} \frac{1}{3} r^3 \sin \theta \Big|_0^4 d\theta =$$

$$\int_0^{\pi/2} \frac{64}{3} \sin \theta d\theta = -\frac{64}{3} \cos \theta \Big|_0^{\pi/2} = -\frac{64}{3} \cos(\pi/2) + \frac{64}{3} \cos(0) = \frac{64}{3}$$



4. Evaluate the integral  $\int_0^2 \int_0^{\sqrt{4-x^2}} \int_0^{\sqrt{x^2+y^2}} \frac{1}{x^2+y^2} dz dy dx$  in cylindrical coordinates. (10 points)

$x=2$   
 $y=\sqrt{4-x^2}$   
 $z=\sqrt{x^2+y^2} \Rightarrow z=r$   
 $x=0$   
 $y=0$   
 $x^2+y^2=r$



$$\int_0^{\pi/2} \int_0^2 \int_0^r \frac{1}{r^2} \cdot r \, dz \, dr \, d\theta =$$

$$\int_0^{\pi/2} \int_0^2 \int_0^r \frac{1}{r} \, dz \, dr \, d\theta = \int_0^{\pi/2} \int_0^2 z \cdot \frac{1}{r} \Big|_0^r \, dr \, d\theta =$$

$$\int_0^{\pi/2} \int_0^2 \frac{r}{r} \, dr \, d\theta = \int_0^{\pi/2} \int_0^2 dr \, d\theta = \int_0^{\pi/2} r \Big|_0^2 \, d\theta = \int_0^{\pi/2} 2 \, d\theta$$

$$= 2\theta \Big|_0^{\pi/2} = \boxed{\pi}$$

5. Set up an iterated integral to find the area of the region bounded by  $2x - 3y = 0$ ,  $x + y = 4$ ,  $y = 0$ . Evaluate the integral to find the area. (10 points)

$$\int_0^{8/5} \int_{\frac{3}{2}y}^{-y+4} 1 \, dx \, dy = \int_0^{8/5} x \Big|_{\frac{3}{2}y}^{-y+4} \, dy =$$


$$\int_0^{8/5} -y+4 - \frac{3}{2}y \, dy$$

$$\int_0^{8/5} -\frac{5}{2}y + 4 \, dy = -\frac{5}{4}y^2 + 4y \Big|_0^{8/5}$$

$$-\frac{5}{4} \left(\frac{8}{5}\right)^2 + 4\left(\frac{8}{5}\right) - 0 =$$

$$-\frac{8}{4} \left(\frac{16}{5}\right) + \frac{32}{5}$$

$$-\frac{16}{5} + \frac{32}{5} = \boxed{\frac{16}{5}}$$



$$\frac{3}{2}y = -y + 4$$

$$3y = -2y + 8$$

$$5y = 8$$

$$y = \frac{8}{5}$$

6. Use a double integral to find the volume of the indicated solid:  $z = x + y, x^2 + y^2 = 1$ , first octant. (10 points)

$$\rightarrow z = x + y \Rightarrow z = r \cos \theta + r \sin \theta$$



$$z = r(\cos \theta + \sin \theta)$$

$$\int_0^{\pi/2} \int_0^1 r(\cos \theta + \sin \theta) r dr d\theta$$

$$\int_0^{\pi/2} \frac{1}{3} r^3 (\cos \theta + \sin \theta) \Big|_0^1 d\theta =$$

$$\frac{1}{3} \int_0^{\pi/2} \cos \theta + \sin \theta d\theta = \frac{1}{3} [\sin \theta - \cos \theta]_0^{\pi/2} =$$

$$\frac{1}{3} [1 - 0 - (0 - 1)] = \boxed{\frac{2}{3}}$$

7. Find the area of the surface given by  $f(x, y) = 4 + 2x + 2y$  over the region  $R$ : rectangle with vertices  $(0,0), (2,0), (2,4), (0,4)$ . (10 points)

$$0 \leq x \leq 2 \quad 0 \leq y \leq 4$$

$$f_x = 2, f_y = 2$$


$$\sqrt{1 + (f_x)^2 + (f_y)^2} = \sqrt{1 + 4 + 4} = \sqrt{9} = 3$$

$$\int_0^2 \int_0^4 3 dy dx =$$

$$\int_0^2 3y \Big|_0^4 dx = \int_0^2 12 dx = 12x \Big|_0^2 = \boxed{24}$$

8. Find the mass and centroid of the center of mass of the lamina bounded by the graphs of the equations  $xy = 4$ ,  $x = 1$ ,  $x = 4$ ,  $y = 0$  with the density  $\rho = ky^2$ . (15 points)

$$y = \frac{4}{x}$$



$$M = \int_1^4 \int_0^{4/x} ky^2 dy dx = \int_1^4 \left. \frac{k}{3} y^3 \right|_0^{4/x} dx =$$

$$\int_1^4 \frac{k \cdot 64}{3x^3} dx = \int_1^4 \frac{k \cdot 64}{3} \cdot x^{-3} dx = -\frac{32k}{3} x^{-2} \Big|_1^4 = -\frac{32k}{3} \cdot \frac{1}{4} + \frac{32k}{3} \cdot \frac{1}{1} = 10k$$

$$M_x = \int_1^4 \int_0^{4/x} ky^3 dy dx = \int_1^4 \left. \frac{k}{4} y^4 \right|_0^{4/x} dx = \int_1^4 \frac{k}{4} \cdot \frac{64}{x^4} dx =$$

$$\int_1^4 64k x^{-4} dx = -\frac{64k}{3} \cdot x^{-3} \Big|_1^4 = -\frac{64k}{3} \cdot \frac{1}{64} + \frac{64k}{3} \cdot \frac{1}{1} = 21k$$

$$M_y = \int_1^4 \int_0^{4/x} kxy^2 dy dx = \int_1^4 \left. \frac{k}{3} xy^3 \right|_0^{4/x} dx = \int_1^4 \frac{k \cdot 64}{3x^3} \cdot x dx =$$

$$\int_1^4 \frac{64k}{3} \cdot x^{-2} dx = -\frac{64k}{3} \cdot \frac{1}{x} \Big|_1^4 = -\frac{64k}{3} \cdot \frac{1}{4} + \frac{64k}{3} \cdot \frac{1}{1} = 16k$$

$$\bar{x} = \frac{M_y}{M} = \frac{16k}{10k} = \frac{16}{10} = \frac{8}{5}$$

$$\bar{y} = \frac{M_x}{M} = \frac{21k}{10k} = \frac{21}{10}$$

$$(\bar{x}, \bar{y}) = \left( \frac{8}{5}, \frac{21}{10} \right)$$

9. Find integrals for  $I_x, I_y, I_z$  for the cube  $Q: \{(x, y, z): 0 \leq x \leq 1, 0 \leq y \leq 1, 0 \leq z \leq 1\}$  with density  $\rho(x, y, z) = kx^2y^2z^2$ . You do not need to integrate them. (10 points)

$$I_x = \int_0^1 \int_0^1 \int_0^1 kx^2y^2z^2(y^2+z^2) dz dy dx$$

$$I_y = \int_0^1 \int_0^1 \int_0^1 kx^2y^2z^2(x^2+z^2) dz dy dx$$

$$I_z = \int_0^1 \int_0^1 \int_0^1 kx^2y^2z^2(x^2+y^2) dz dy dx$$

10. Find the volume of the solid between the cone  $z = \sqrt{x^2 + y^2}$  and the sphere  $x^2 + y^2 + z^2 = 12$ . (10 points)

$$\rho \cos \varphi = \rho \sin \varphi \Rightarrow \varphi = \pi/4$$

$$\int_0^{\pi/4} \int_0^{2\pi} \int_0^{\sqrt{12}} \rho^2 \sin \varphi d\rho d\theta d\varphi$$

$$\int_0^{\pi/4} \int_0^{2\pi} \left. \frac{\rho^3}{3} \right|_0^{\sqrt{12}} \sin \varphi d\theta d\varphi = \int_0^{\pi/4} \int_0^{2\pi} \frac{4\sqrt{12}}{3} \sin \varphi d\theta d\varphi$$

$$\int_0^{\pi/4} 4\sqrt{12} \sin \varphi \theta \Big|_0^{2\pi} d\varphi = \int_0^{\pi/4} 8\pi\sqrt{12} \sin \varphi d\varphi = -8\pi\sqrt{12} \cos \varphi \Big|_0^{\pi/4}$$

$$= -8\pi\sqrt{12} \cdot \left( \frac{1}{\sqrt{2}} - 1 \right) = \boxed{\frac{16\pi\sqrt{3}(\sqrt{2}-1)}{\sqrt{2}}}$$

